





Country Profile: Malaysia

Last Updated: 07 December 2023 **Region:** South-Eastern Asia



Identified policies and legal sources related to abortion:

- Reproductive Health Act
- ✓ General Medical Health Act
- Constitution
- ✓ Criminal / Penal Code
 - Civil Code
 - Ministerial Order / Decree
 - Case Law
- ✓ Health Regulation / Clinical Guidelines
- ✓ EML / Registered List
 - Medical Ethics Code
 - **Document Relating to Funding**
- Abortion Specific Law
- Law on Medical Practicioners
- Law on Health Care Services
- Other

Related Documents

From General Medical Health Act:

• Private Healthcare Facilities and Services Act, 1998

From Criminal / Penal Code:

• Penal Code, 2006

From Health Regulation / Clinical Guidelines:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012

From EML / Registered List:

• National Essential Medicines List, 2023



Concluding Observations:

• HEALTH



Persons who can be sanctioned:

- ✓ A woman or girl can be sanctioned
- ✓ Providers can be sanctioned ✓ A person who assists can be sanctioned



List of ratified human rights treaties:

CERD

CCPR

Xst OP

2nd OP

CESCR

CESCR-OP

CAT

CAT-OP

CEDAW

CEDAW-OP

✓ CRC

✓ CRC:OPSC

✓ CRC:OPAC CRC:OPIC

CMW

✓ CRPD * CRPD-OP

CED **

Maputo Protocol

 ψ Download data

Abortion at the woman's request



Legal Ground and Gestational Limit

Economic or social

No

Related documents:

• Penal Code, 2006 (page 126)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Health grounds shall reflect WHO's definitions of health, which entails a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity. Abortion Care Guideline § 2.2.2.

↓ Source document: WHO Abortion Care Guideline (page 16)

Foetal impairment

No

Related documents:

• Penal Code, 2006 (page 126)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Grounds-based approaches to restricting access to abortion should be revised in favour of making abortion available on the request of the woman, girl or other pregnant person. The Abortion Care Guideline recommends against laws and other regulations that restrict abortion by grounds. The guideline recommends abortion be available on the request of the woman, girl or other pregnant person.

Until they are replaced with abortion on request, any existing grounds should be formulated and applied in a manner consistent with international human rights law. This requires that abortion is available when carrying a pregnancy to term would cause the woman, girl or other pregnant person substantial pain or suffering, including but not limited to situations where the pregnancy is not viable. Grounds-based approaches that require fetal impairments to be fatal for abortion to be lawful frustrate providers and leave women no choice but to continue with pregnancy. Being required to continue with a pregnancy that causes significant distress violates numerous human rights. Abortion Care Guideline § 2.2.2.

↓ Source document: WHO Abortion Care Guideline (page 64)

Rape

No

Related documents:

• Penal Code, 2006 (page 126)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Grounds-based approaches to restricting access to abortion should be revised in favour of making abortion available on the request of the woman, girl or other pregnant person. The Abortion Care Guideline recommends against laws and other regulations that restrict abortion by grounds. The guideline recommends abortion be available on the request of the woman, girl or other pregnant person.

Until they are replaced with abortion on request, any existing grounds should be formulated and applied in a manner consistent with international human rights law. This requires that abortion is available when carrying a pregnancy to term would cause the woman, girl or other pregnant person substantial pain or suffering, including but not limited to situations where the pregnancy is the result of rape or incest Abortion Care Guideline § 2.2.2.

Source document: WHO Abortion Care Guideline (page 64)

Incest

No

Related documents:

• Penal Code, 2006 (page 126)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Grounds-based approaches to restricting access to abortion should be revised in favour of making abortion available on the request of the woman, girl or other pregnant person. The Abortion Care Guideline recommends against laws and other regulations that restrict abortion by grounds. The guideline recommends abortion be available on the request of the woman, girl or other pregnant person.

Until they are replaced with abortion on request, any existing grounds should be formulated and applied in a manner consistent with international human rights law. This requires that abortion is available when carrying a pregnancy to term would cause the woman, girl or other pregnant person substantial pain or suffering, including but not limited to situations where the pregnancy is the result of rape or incest. Abortion Care Guideline § 2.2.2.

→ Source document: WHO Abortion Care Guideline (page 64)

Intellectual or cognitive disability of the woman

No

Related documents:

• Penal Code, 2006 (page 126)

Mental health

Yes

Related documents:

• Penal Code, 2006 (page 126)

Gestational limit

Weeks: 22

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:6)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Health grounds shall reflect WHO's definitions of health, which entails a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity. Abortion Care Guideline § 2.2.2.

→ Source document: WHO Abortion Care Guideline (page 16)

Laws or policies that impose time limits on the length of pregnancy may have negative consequences for women, including forcing them to seek clandestine abortions and suffer social inequities. Safe Abortion Guidelines, § 4.2.1.7.

→ Source document: WHO Abortion Care Guideline (page 103)

Physical health	Yes
	Related documents:
	• Penal Code, 2006 (page 126)
	Gestational limit
	Weeks: 22
	• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:6)
	WHO Guidance
	The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.
	Health grounds shall reflect WHO's definitions of health, which entails a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity. Abortion Care Guideline § 2.2.2.
	→ Source document: WHO Abortion Care Guideline (page 16)
	Laws or policies that impose time limits on the length of pregnancy may have negative consequences for women, including forcing them to seek clandestine abortions and suffer social
	inequities. Safe Abortion Guidelines, § 4.2.1.7.
	→ Source document: WHO Abortion Care Guideline (page 103)
Health	No
	Related documents:
	• Penal Code, 2006 (page 126)
	WHO Guidance
	The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.
	Health grounds shall reflect WHO's definitions of health, which entails a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity. Abortion
	Care Guideline § 2.2.2.
	↓ Source document : WHO Abortion Care Guideline (page 16)
	Vac
Life	Yes
	Related documents: • Penal Code, 2006 (page 126)
	Tenan code, 2000 (page 120)
	Gestational limit
	Weeks: 22
	• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:6)
	WHO Guidance
	The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.
	Grounds-based approaches to restricting access to abortion should be revised in favour of making abortion available on the request of the woman, girl or other pregnant person. The Abortion
	Care Guideline recommends against laws and other regulations that restrict abortion by grounds. The guideline recommends abortion be available on the request of the woman, girl or other pregnant person.
	Until they are replaced with abortion on request, any existing grounds should be formulated and applied in a manner consistent with international human rights law. This requires that abortion is available where the life and health of the woman, girl or other pregnant person is at risk. Abortion Care Guideline § 2.2.2.
	→ Source document: WHO Abortion Care Guideline (page 64)
	Laws or policies that impose time limits on the length of pregnancy may have negative consequences for women, including forcing them to seek clandestine abortions and suffer social
	inequities. Safe Abortion Guidelines, § 4.2.1.7.
	→ Source document : WHO Abortion Care Guideline (page 103)

Other

Authorization of health professional(s)

Yes

Related documents:

- Penal Code, 2006 (page 126)
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)

Number and cadre of health-care professional authorizations required

1

Registered Medical Practitioner

While by law, only one medical registered practitioner is required to

assess if a termination of pregnancy is, it is suggested that in a Government Hospital setting, two doctors, one of whom is a specialist, should concur on the termination of Pregnancy. For mental health reasons, an opinion from a psychologist or psychiatrist is not needed unless it is deemed necessary by the attending doctor i.e. because of severe depression or suicidal risk.

- Penal Code, 2006 (page 126)
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 7)
- Penal Code, 2006 (page 126)
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 7)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Third-party authorization requirements are incompatible with international human rights law, which provides that States may not restrict women's access to health services on the ground that they do not have the authorization of husbands, partners, parents or health authorities, because they are unmarried, or because they are women. The Abortion Care Guideline recommends that abortion be available on the request of the woman, girl or other pregnant person without the authorization of any other individual, body or institution. Abortion Care Guideline § 3.3.2.

→ Source document: WHO Abortion Care Guideline (page 81)

Authorization in specially licensed facilities only



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

To establish an enabling environment, there is a need for abortion care to be integrated into the health system across all levels (including primary, secondary and tertiary) – and supported in the community – to allow for expansion of health worker roles, including self-management approaches. To ensure both access to abortion and achievement of Universal Health Coverage (UHC), abortion must be centred within primary health care (PHC), which itself is fully integrated within the health system, facilitating referral pathways for higher-level care when needed. Abortion Care Guideline § 1.4.1.

→ Source document: WHO Abortion Care Guideline (page 52)

Judicial authorization for minors



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends that abortion be available on the request of the woman, girl or other pregnant person without the authorization of any other individual, body or institution. Abortion Care Guideline § 3.3.2.

→ Source document: WHO Abortion Care Guideline (page 81)

Judicial authorization in cases of rape

not applicable



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

There shall be no procedural requirements to "prove" or "establish" satisfaction of grounds, such as requiring judicial orders or police reports in cases of rape or sexual assault (for sources to support this information). These restrictions subject the individual to unnecessary trauma, may put them at increased risk from the perpetrator, and may cause women to resort to unsafe abortion.

The Abortion Care Guideline recommends abortion be available on the request of the woman, girl or other pregnant person. Abortion Care Guideline § 2.2.2.

→ Source document: WHO Abortion Care Guideline (page 64)

Police report required in case of rape

not applicable



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

There shall be no procedural requirements to "prove" or "establish" satisfaction of grounds, such as requiring judicial orders or police reports in cases of rape or sexual assault (for sources to support this information). These restrictions subject the individual to unnecessary trauma, may put them at increased risk from the perpetrator, and may cause women to resort to unsafe abortion.

The Abortion Care Guideline recommends abortion be available on the request of the woman, girl or other pregnant person. Abortion Care Guideline § 2.2.2.

↓ Source document: WHO Abortion Care Guideline (page 64)

Parental consent required for minors

Yes

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:9)

Can another adult consent in place of a parent?

VΔC

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:9)

Age where consent not needed

18

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:9)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

While parental or partner involvement in abortion decision-making can support and assist women, girls or other pregnant persons, this must be based on the values and preferences of the person availing of abortion and not imposed by third-party authorization requirements. Third-party authorization requirements are incompatible with international human rights law, which provides that States may not restrict women's access to health services on the ground that they do not have the authorization of husbands, partners, parents or health authorities, because they are unmarried, or because they are women. The Abortion Care Guideline recommends that abortion be available on the request of the woman, girl or other pregnant person without the authorization of any other individual, body or institution. Abortion Care Guideline § 3.3.2.

↓ Source document: WHO Abortion Care Guideline (page 81)

Spousal consent

No

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:9)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

While parental or partner involvement in abortion decision-making can support and assist women, girls or other pregnant persons, this must be based on the values and preferences of the person availing of abortion and not imposed by third-party authorization requirements. Third-party authorization requirements are incompatible with international human rights law, which provides that States may not restrict women's access to health services on the ground that they do not have the authorization of husbands, partners, parents or health authorities, because they are unmarried, or because they are women. The Abortion Care Guideline recommends that abortion be available on the request of the woman, girl or other pregnant person without the authorization of any other individual, body or institution. Abortion Care Guideline § 3.3.2.

→ Source document: WHO Abortion Care Guideline (page 81)



Additional notes

Muslim women require husband's consent as per Fatwa, this is not a legal requirement otherwise.

Ultrasound images or listen to foetal heartbeat required

No

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:8)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The right to refuse information, including the right to refuse viewing ultrasound images, must be respected. The Abortion Care Guideline recommends against the use of ultrasound scanning as a prerequisite for providing abortion services for both medical and surgical abortion. Abortion Care Guideline § 3.3.5.

↓ Source document: WHO Abortion Care Guideline (page 85)

Compulsory counselling

No

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:23)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

While counselling should be made available and accessible, it should always be voluntary for women to choose whether or not they want to receive it. The right to refuse counselling when offered must be respected. Where provided, counselling must be available to individuals in a way that respects privacy and confidentiality.

Counselling should be person-centred and may need to be tailored according to the needs of the individual; young people, survivors of sexual and gender-based violence or members of marginalized groups may have different information or counselling requirements.

The content of and approach to counselling will need to be adjusted depending on the reason for seeking abortion services. Therefore, it is important for the counsellor to be aware of and sensitive to the individual's situation and needs. Abortion Care Guideline § 3.2.2.

↓ Source document: WHO Abortion Care Guideline (page 77)

Compulsory waiting period

Yes

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:8)

Waiting period

After the first counseling by the health care professional 48 HOURS



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Mandatory waiting periods delay access to abortion, sometimes to the extent that women's access to abortion or choice of abortion method is restricted. The Abortion Care Guideline recommends against mandatory waiting periods for abortion. Abortion Care Guideline § 3.3.1.

Source document: WHO Abortion Care Guideline (page 79)

Mandatory HIV screening test

No

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:8)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Regulatory, policy and programmatic barriers – as well as barriers in practice – that hinder access to and timely provision of quality abortion care should be removed. Abortion Care Guideline § Box 2.1.

Source document: WHO Abortion Care Guideline (page 59)



Additional notes

HIV and hepatitis testing should only be performed if indicated in the light of clinical features, individual risk factors, or local prevalence.

Other mandatory STI screening tests

No

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:8)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Regulatory, policy and programmatic barriers – as well as barriers in practice – that hinder access to and timely provision of quality abortion care should be removed. Abortion Care Guideline § Box 2.1.

→ Source document: WHO Abortion Care Guideline (page 59)



Additional notes

HIV and hepatitis testing should only be performed if indicated in the light of clinical features, individual risk factors, or local prevalence.

Prohibition of sexselective abortion



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

• Penal Code, 2006



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

In situations where abortion is restricted for sex selection purposes, terminating a pregnancy for this reason is likely to involve an unsafe procedure carrying high risks. Any policies or guidelines on the use of technology in obstetric and fetal medicine should take into account the need to ensure women's access to safe abortion and other services - efforts to manage or limit sex selection should also not hamper or limit access to safe abortion services. Preventing gender-biased sex selection: an interagency statement.

↓ Source document: Preventing Gender-Biased Sex Selection (page 17)

Restrictions on information provided to the public

No data found



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Dissemination of misinformation, withholding of information and censorship should be prohibited.

Information should be accessible and understandable, including formats catering to low-literacy and differently abled populations. Different modalities exist for the provision of information on abortion, e.g. remote access via hotlines and telemedicine, and through approaches such as harm reduction and community-based outreach, as well as in-person interactions with health workers. Abortion Care Guideline § 3.2.1.

Source document: WHO Abortion Care Guideline (page 74)

Restrictions on methods to detect sex of the foetus



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

A woman is entitled to know the status of her pregnancy and to act on this information. Prenatal tests and other medical diagnostic services cannot legally be refused because the woman may decide to terminate her pregnancy. Safe Abortion Guidelines § 4.2.1.4.

◆ Source document: WHO Abortion Care Guideline (page 103)

Other

Consent for cognitively impaired women may be obtained from parent/guardian or doctors caring for her

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:9)

Clinical and Service-delivery Aspects of Abortion Care

National guidelines for induced abortion

Yes, guidelines issued by the government

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page A)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

National standards and guidelines for abortion care should be evidence based and periodically updated and should provide the necessary guidance to achieve equal access to comprehensive abortion care. Leadership should also promote evidence-based SRH services according to these standards and guidelines. Abortion Care Guideline § 1.3.3.

↓ Source document: WHO Abortion Care Guideline (page 50)

Methods allowed

Vacuum aspiration

Yes (7-19 WEEKS)

Manual vacuum aspiration is permitted below ten weeks of gestation. Suction evacuation (vacuum aspiration/suction curettage) is permitted between seven and 14 weeks, and surgical evacuation is permitted up to 19 weeks.

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:11)

Dilatation and evacuation

Yes (15-20 WEEKS)

Dilation and evacuation is to be done by a specialist practitioner. It is not the method of

choice since it is associated with complications such as bleeding and

cervical injury.

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:20)

Combination mifepristone-misoprostol

Yes (14-22 WEEKS)

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:11)

Misoprostol only

Yes (5-9, 14-22 WEEKS)

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:11)

Other (where provided)

Gemeprostor or methotrexate with misoprostol Extra-amniotic saline infiltration is also listed, but with a note regarding limited safety/efficacy data

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:11)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Vacuum aspiration is recommended for surgical abortions at or under 14 weeks to be provided by traditional and complementary medicine professionals, nurses, midwives, associate/advanced associate clinicians, generalist medical practitioners and specialist medical practitioners.

The Abortion Care Guideline recommends against the practice of dilatation and sharp curettage (D&C), including for sharp curette checks (i.e. to "complete" the abortion) following vacuum aspiration. Abortion Care Guideline § 3.4.1.

↓ Source document: WHO Abortion Care Guideline (page 101)

Dilation and evacuation (D&E) is recommended for surgical abortions at or over 14 weeks to be provided by generalist medical practitioners and specialist medical practitioners. Vacuum aspiration can be used during a D&E. Abortion Care Guideline § 3.4.1.

→ Source document: WHO Abortion Care Guideline (page 103)

The recommended method for medical abortion is mifepristone followed by misoprostol (regimen differs by gestational age). Abortion Care Guideline § 3.4.2.

→ Source document: WHO Abortion Care Guideline (page 106)

The Abortion Care Guideline recommends the use of misoprostol alone, with a regime that differs by gestational age. Evidence demonstrates that the use of combination mifepristone plus misoprostol is more effective than misoprostol alone. Abortion Care Guideline § 3.4.2.

✓ Source document: WHO Abortion Care Guideline (page 106)

Country recognized approval (mifepristone / mifemisoprostol)

No

Related documents:

• National Essential Medicines List, 2023 (page 1)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Mifepristone and misoprostol should be listed in relevant national EMLs (NEMLs) or their equivalent and should be included in the relevant clinical care/service delivery guidelines.

Inclusion in the NEML is one important component of ensuring that quality medicines are available.

For induced abortion, Mifepristone (200 mg) and misoprostol (200 μ g), are recommended in the WHO EML. The EML specifically mentions the following co-packaged formulation: 1 tablet mifepristone (200 mg) + 4 tablets misoprostol (200 μ g).

Restrictions on prescribing authority for some categories of health workers may need to be modified or other mechanisms put in place to make the medicines available for these health workers within the regulatory framework of the health system. Abortion Care Guideline § 1.4.4.

→ Source document: WHO Abortion Care Guideline (page 55)

Mifepristone and misoprostol should be listed in relevant national EMLs (NEMLs) or their equivalent and should be included in the relevant clinical care/service delivery guidelines.

Inclusion in the NEML is one important component of ensuring that quality medicines are available.

For induced abortion, Mifepristone (200 mg) and misoprostol (200 μ g), are recommended in the WHO EML. The EML specifically mentions the following co-packaged formulation: 1 tablet mifepristone (200 mg) + 4 tablets misoprostol (200 μ g).

Restrictions on prescribing authority for some categories of health workers may need to be modified or other mechanisms put in place to make the medicines available for these health workers within the regulatory framework of the health system. Abortion Care Guideline § 1.4.4.

↓ Source document: WHO Abortion Care Guideline (page 55)

Country recognized approval (misoprostol)

Yes, for gynaecological indications

Related documents:

• National Essential Medicines List, 2023 (page 30)

Misoprostol allowed to be sold or distributed by pharmacies or drug stores



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

• National Essential Medicines List, 2023



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Mifepristone and misoprostol should be listed in relevant national EMLs (NEMLs) or their equivalent and should be included in the relevant clinical care/service delivery guidelines.

Inclusion in the NEML is one important component of ensuring that quality medicines are available.

For induced abortion, Mifepristone (200 mg) and misoprostol (200 μ g) are recommended in the WHO EML. The EML specifically mentions the following co-packaged formulation: 1 tablet mifepristone (200 mg) + 4 tablets misoprostol (200 μ g).

Restrictions on prescribing authority for some categories of health workers may need to be modified or other mechanisms put in place to make the medicines available for these health workers within the regulatory framework of the health system. Abortion Care Guideline § 1.4.4.

↓ Source document: WHO Abortion Care Guideline (page 55)

Where can abortion services be provided

Related documents:

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)
- Private Healthcare Facilities and Services Act, 1998 (page 1)

Primary health-care centres

Not specified

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998

Secondary (district-level) health-care facilities

Yes

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)
- Private Healthcare Facilities and Services Act, 1998 (page 1)

Specialized abortion care public facilities

Not specified

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998

Private health-care centres or clinics

Yes

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)
- Private Healthcare Facilities and Services Act, 1998 (page 1)

NGO health-care centres or clinics

Not specified

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Where it is lawful, abortion must be accessible in practice. This requires both ensuring that health-care facilities, commodities and services are accessible (including sufficient providers), and that law and policy on abortion is formulated, interpreted and applied in a way that is compatible with human rights. Abortion Care Guideline § 1.3.1.

→ Source document: WHO Abortion Care Guideline (page 48)



Additional notes

The Guidelines on Termination of Pregnancy state: "Patients can be seen at any private or public health institution but the procedure should only be done in a setting with Gynaecologist (Specialist) support (this is also in accordance with the Private Health Care Act). This is to ensure that the procedure can be done properly and if complications should occur, these complications can be picked up quickly."

National guidelines for post-abortion care

Yes, guidelines issued by the government

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page A)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

National standards and guidelines for abortion care should be evidence based and periodically updated and should provide the necessary guidance to achieve equal access to comprehensive abortion care. Leadership should also promote evidence-based SRH services according to these standards and guidelines. Abortion Care Guideline § 1.3.3.

→ Source document: WHO Abortion Care Guideline (page 50)

Where can post abortion care services be provided

Primary health-care centres

Not specified

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998

Secondary (district-level) health-care facilities

Yes

The Guidelines on Termination of Pregnancy state: "Patients can be seen at any private or public health institution but the procedure should only be done in a setting with Gynaecologist (Specialist) support (this is also in accordance with the Private Health Care Act). This is to ensure that the procedure can be done properly and if complications should occur, these complications can be picked up quickly."

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)
- Private Healthcare Facilities and Services Act, 1998 (page 1)

Specialized abortion care public facilities

Not specified

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998

Private health-care centres or clinics

Yes

The Guidelines on Termination of Pregnancy state: "Patients can be seen at any private or public health institution but the procedure should only be done in a setting with Gynaecologist (Specialist) support (this is also in accordance with the Private Health Care Act). This is to ensure that the procedure can be done properly and if complications should occur, these complications can be picked up quickly."

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)
- Private Healthcare Facilities and Services Act, 1998 (page 1)

NGO health-care centres or clinics

Not specified

The Guidelines on Termination of Pregnancy state: "Patients can be seen at any private or public health institution but the procedure should only be done in a setting with Gynaecologist (Specialist) support (this is also in accordance with the Private Health Care Act). This is to ensure that the procedure can be done properly and if complications should occur, these complications can be picked up quickly."

- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012
- Private Healthcare Facilities and Services Act, 1998
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)
- Private Healthcare Facilities and Services Act, 1998 (page 1)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends the option of telemedicine as an alternative to in-person interactions with the health worker to deliver medical abortion services in whole or in part. Telemedicine services should include referrals (based on the woman's location) for medicines (abortion and pain control medicines), any abortion care or post-abortion follow-up required (including for emergency care if needed), and for post-abortion contraceptive services. Abortion Care Guideline § 3.6.1.

↓ Source document: WHO Abortion Care Guideline (page 133)

Contraception included in post-abortion care

Yes

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:13)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

All contraceptive options may be considered after an abortion. For individuals undergoing surgical abortion and wishing to use contraception, Abortion Care Guideline recommends the option of initiating the contraception at the time of surgical abortion. For individuals undergoing medical abortion, for those who choose to use hormonal contraception, the Abortion Care Guideline suggests that they be given the option of starting hormonal contraception immediately after the first pill of the medical abortion regimen. For those who choose to have an IUD inserted, Abortion Care Guideline suggests IUD placement at the time that success of the abortion procedure is determined. Abortion Care Guideline § 3.5.4.

↓ Source document: WHO Abortion Care Guideline (page 126)

Insurance to offset end user costs



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012

Other (if applicable)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Where user fees are charged for abortion, this should be based on careful consideration of ability to pay, and fee waivers should be available for those who are facing financial hardship and adolescent abortion seekers. As far as possible, abortion services and supplies should be mandated for coverage under insurance plans as inability to pay is not an acceptable reason to deny or delay abortion care. Furthermore, having transparent procedures in all health-care facilities can ensure that informal charges are not imposed by staff. Abortion Care Guideline § 1.4.2.

→ Source document: WHO Abortion Care Guideline (page 53)

Who can provide abortion services

Related documents:

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:15)

Nurse

Not specified

• Penal Code, 2006

Midwife/nurse-midwife

Not specified

• Penal Code, 2006

Doctor (specialty not specified)

Not specified

• Penal Code, 2006

Specialist doctor, including OB/GYN

Not specified

• Penal Code, 2006

Other (if applicable)

Medical Practitioner Registered under the Medical Act 1971

• Penal Code, 2006 (page 126)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends against regulation on who can provide and manage abortion that is inconsistent with WHO guidance. Abortion Care Guideline § 3.3.8.

Source document: WHO Abortion Care Guideline (page 97)

Extra facility/provider requirements for delivery of abortion services

Referral linkages to a higher-level facility

Not specified

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012

Availability of a specialist doctor, including OB/GYN

Yes

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012 (page 1:7)

Minimum number of beds

Not specified

• Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012

Other (if applicable)



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

There is no single recommended approach to providing abortion services. The choice of specific health worker(s) (from among the recommended options) or management by the individual themself, and the location of service provision (from among recommended options) will depend on the values and preferences of the woman, girl or other pregnant person, available resources, and the national and local context. A plurality of service-delivery approaches can co-exist within any given context. Given that service-delivery approaches can be diverse, it is important to ensure that for the individual seeking care, the range of service-delivery options taken together will provide access to scientifically accurate, understandable information at all stages; access to quality-assured medicines (including those for pain management); back-up referral support if desired or needed; linkages to an appropriate choice of contraceptive services for those who want post-abortion contraception. Best Practice Statement 49 on service delivery. Abortion Care Guideline § 3.6.1.

→ Source document: WHO Abortion Care Guideline (page 132)



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends that access to and continuity of comprehensive abortion care be protected against barriers created by conscientious objection. It is critical that States ensure compliance with regulations and design/organize health systems to ensure access to and continuity of quality abortion care. If it proves impossible to regulate conscientious objection in a way that respects, protects and fulfils abortion seekers' rights, conscientious objection in abortion provision may become indefensible. Abortion Care Guideline § 3.3.9.

↓ Source document: WHO Abortion Care Guideline (page 98)



Additional notes

The Professional Code of Conduct of the Malaysian Medical Council states in Section 1.2.1: "It should be remembered that a practitioner suffers no loss of dignity or prestige in referring a patient to a colleague whose opinion could contribute to the better care of the patient." The Code of Conduct can be accessed at: http://www.dermaorgan.gov.my/pdf/Code%20of%20Professional%20Conduct%20-%20Malaysian%20Medical%20Council.pdf

Private sector providers



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends that access to and continuity of comprehensive abortion care be protected against barriers created by conscientious objection. It is critical that States ensure compliance with regulations and design/organize health systems to ensure access to and continuity of quality abortion care. If it proves impossible to regulate conscientious objection in a way that respects, protects and fulfils abortion seekers' rights, conscientious objection in abortion provision may become indefensible. Abortion Care Guideline § 3.3.9.

↓ Source document: WHO Abortion Care Guideline (page 98)



Additional notes

The Professional Code of Conduct of the Malaysian Medical Council states in Section 1.2.1: "It should be remembered that a practitioner suffers no loss of dignity or prestige in referring a patient to a colleague whose opinion could contribute to the better care of the patient." The Code of Conduct can be accessed at: http://www.dermaorgan.gov.my/pdf/Code%20of%20Professional%20Conduct%20-%20Malaysian%20Medical%20Council.pdf

Provider type not specified



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends that access to and continuity of comprehensive abortion care be protected against barriers created by conscientious objection. It is critical that States ensure compliance with regulations and design/organize health systems to ensure access to and continuity of quality abortion care. If it proves impossible to regulate conscientious objection in a way that respects, protects and fulfils abortion seekers' rights, conscientious objection in abortion provision may become indefensible. Abortion Care Guideline § 3.3.9.

→ Source document: WHO Abortion Care Guideline (page 98)



Additional notes

The Professional Code of Conduct of the Malaysian Medical Council states in Section 1.2.1: "It should be remembered that a practitioner suffers no loss of dignity or prestige in referring a patient to a colleague whose opinion could contribute to the better care of the patient." The Code of Conduct can be accessed at: http://www.dermaorgan.gov.my/pdf/Code%20of%20Professional%20Conduct%20-%20Malaysian%20Medical%20Council.pdf

Neither Type of Provider Permitted



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

The Abortion Care Guideline recommends that access to and continuity of comprehensive abortion care be protected against barriers created by conscientious objection. It is critical that States ensure compliance with regulations and design/organize health systems to ensure access to and continuity of quality abortion care. If it proves impossible to regulate conscientious objection in a way that respects, protects and fulfils abortion seekers' rights, conscientious objection in abortion provision may become indefensible. Abortion Care Guideline § 3.3.9.

↓ Source document: WHO Abortion Care Guideline (page 98)



Additional notes

The Professional Code of Conduct of the Malaysian Medical Council states in Section 1.2.1: "It should be remembered that a practitioner suffers no loss of dignity or prestige in referring a patient to a colleague whose opinion could contribute to the better care of the patient." The Code of Conduct can be accessed at: http://www.dermaorgan.gov.my/pdf/Code%20of%20Professional%20Conduct%20-%20Malaysian%20Medical%20Council.pdf

Public facilities



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Where it is lawful, abortion must be accessible in practice. This requires both ensuring that health-care facilities, commodities and services are accessible (including sufficient providers), and that law and policy on abortion is formulated, interpreted and applied in a way that is compatible with human rights. Abortion Care Guideline § 1.3.1.

↓ Source document: WHO Abortion Care Guideline (page 48)

Private facilities



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Where it is lawful, abortion must be accessible in practice. This requires both ensuring that health-care facilities, commodities and services are accessible (including sufficient providers), and that law and policy on abortion is formulated, interpreted and applied in a way that is compatible with human rights. Abortion Care Guideline § 1.3.1.

→ Source document: WHO Abortion Care Guideline (page 48)

Facility type not specified



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Where it is lawful, abortion must be accessible in practice. This requires both ensuring that health-care facilities, commodities and services are accessible (including sufficient providers), and that law and policy on abortion is formulated, interpreted and applied in a way that is compatible with human rights. Abortion Care Guideline § 1.3.1.

→ Source document: WHO Abortion Care Guideline (page 48)

Neither Type of Facility Permitted



Not specified

When there is no explicit reference to an issue covered in the questionnaire in the relevant document(s), this is noted and no interpretation was made.

Related documents:

- Penal Code, 2006
- Guideline on Termination of Pregnancy for Hospitals in the Ministry of Health, 2012

5.2.2 Proportion of women and girls aged 15 years and older subjected to sexual violence by persons other than an intimate partner in the

5.3.1 Proportion of women aged 20-24 years who were married or in a union before age 15 and before age 18



WHO Guidance

The following descriptions were extracted from WHO Abortion Care Guideline. Where there is a specific Recommendation, this is stated. Otherwise, these are excerpts.

Where it is lawful, abortion must be accessible in practice. This requires both ensuring that health-care facilities, commodities and services are accessible (including sufficient providers), and that law and policy on abortion is formulated, interpreted and applied in a way that is compatible with human rights. Abortion Care Guideline § 1.3.1.

→ Source document: WHO Abortion Care Guideline (page 48)

Indicators

Country specific information related to sexual and reproductive health indications. As data for the Sustainable Development Goal (SDG) indicators related to sexual and reproductive health become available, these will be provided, through periodic updates.

Goal 1. End poverty in all its forms everywhere

previous 12 months, by age and place of occurrence

1.1.1 Proportion of population below the international poverty line, by sex, age, employment status and geographical location (urban/rural)	No data
1.3.1 Proportion of population covered by social protection floors/systems, by sex, distinguishing children, unemployed persons, older persons, persons with disabilities, pregnant women, newborns, work-injury victims and the poor and the vulnerable	No data
La.2 Proportion of total government spending on essential services (education, health and social protection)	No data
Goal 3. Ensure healthy lives and promote well-being for all at all ages	
3.1.1 Maternal mortality ratio	29 (2017)
3.1.2 Proportion of births attended by skilled health personnel	No data
3.7.1 Proportion of women of reproductive age (aged 15-49 years) who have their need for family planning satisfied with modern methods	No data
3.7.2 Adolescent birth rate (aged 10-14 years; aged 15-19 years) per 1,000 women in that age group	13.4 (2015-2020)
3.8.2 Number of people covered by health insurance or a public health system per 1,000 population	No data
3.c.1 Health worker density and distribution	No data
Goal 4. Ensure inclusive and equitable quality education and promote lifelong learning opportunities for all	
4.1.1 Proportion of children and young people: (a) in grades 2/3; (b) at the end of primary; and (c) at the end of lower secondary achieving at east a minimum proficiency level in (i) reading and (ii) mathematics, by sex	No data
Goal 5. Achieve gender equality and empower all women and girls	
5.1.1 Whether or not legal frameworks are in place to promote, enforce and monitor equality and nondiscrimination on the basis of sex	No data
5.2.1 Proportion of ever-partnered women and girls aged 15 years and older subjected to physical, sexual or psychological violence by a current or former intimate partner in the previous 12 months, by form of violence and by age	No data

No data

No data

3.2 Proportion of girls and women aged 15-49 years who have undergone female genital mutilation/cutting, by age	No data
5.1 Proportion of women aged 15-49 years who make their own informed decisions regarding sexual relations, contraceptive use and productive health care	No data
5.2 Number of countries with laws and regulations that guarantee women aged 15- 49 years access to sexual and reproductive health care, brmation and education	No data
.1 (a) Proportion of total agricultural population with ownership or secure rights over agricultural land, by sex; and (b) share of women ong owners or rights-bearers of agricultural land, by type of tenure	No data
.1 Proportion of individuals who own a mobile telephone, by sex	No data
al 8. Promote sustained, inclusive and sustainable economic growth, full and productive employment and decent work for all	
.2 Unemployment rate, by sex, age and persons with disabilities	No data
al 10. Reduce inequality within and among countries	
2.1 Proportion of people living below 50 per cent of median income, by age, sex and persons with disabilities	No data
3.1 Proportion of the population reporting having personally felt discriminated against or harassed within the previous 12 months on the sis of a ground of discrimination prohibited under international human rights law	No data
al 16. Promote peaceful and inclusive societies for sustainable development, provide access to justice for all and build effective, account stitutions at all levels	able and inclusive
1.3 Proportion of population subjected to physical, psychological or sexual violence in the previous 12 months	No data
2.2 Number of victims of human trafficking per 100,000 population, by sex, age and form of exploitation	No data
2.3 Proportion of young women and men aged 1829 years who experienced sexual violence by age 18	No data
3.1 Proportion of victims of violence in the previous 12 months who reported their victimization to competent authorities or other officially ognized conflict resolution mechanisms	No data
5.1 Proportion of persons who had at least one contact with a public official and who paid a bribe to a public official, or were asked for a pe by those public officials, during the previous 12 months	No data
6.1 Primary government expenditures as a proportion of original approved budget, by sector (or by budget codes or similar)	No data
6.2 Proportion of the population satisfied with their last experience of public services	No data
7.1 Proportions of positions (by sex, age, persons with disabilities and population groups) in public institutions (national and local islatures, public service, and judiciary) compared to national distributions	No data
9.1 Proportion of children under 5 years of age whose births have been registered with a civil authority, by age	No data
10.1 Number of verified cases of killing, kidnapping, enforced disappearance, arbitrary detention and torture of journalists, associated dia personnel, trade unionists and human rights advocates in the previous 12 months	No data
b.1 Proportion of population reporting having personally felt discriminated against or harassed in the previous 12 months on the basis of a und of discrimination prohibited under international human rights law	No data
al 17. Strengthen the means of implementation and revitalize the Global Partnership for Sustainable Development	

No data

17.8.1 Proportion of individuals using the Internet

Percentage of married women with unmet need for family planning No data Percentage of births attended by trained health professional 99.4 (2015) Percentage of women aged 20-24 who gave birth before age 18 No data Total fertility rate **2.001** (2018) Legal marital age for women, with parental consent **16** (2009-2017) Legal marital age for women, without parental consent No data Gender Inequalities Index (Value) **0.29** (2017) Gender Inequalities Index (Rank) **62** (2017) Mandatory paid maternity leave **no** (2020) Median age **30.3** (2020) Population, urban (%) **76.036** (2018) Percentage of secondary school completion rate for girls **0.91** (2013) Gender parity in secondary education **1.084** (2018) Percentage of women in non-agricultural employment **39.2** (2013) Proportion of seats in parliament held by women **13.1** (2017)

1.06 (2018)

Additional Reproductive Health Indicators

Sex ratio at birth (male to female births)